

IntechOpen

IntechOpen Series
Sustainable Development, Volume 3

Forest Degradation Under Global Change

Edited by Pavel Samec



Forest Degradation Under Global Change
<http://dx.doi.org/10.5772/intechopen.100671>
Edited by Pavel Samec

Contributors

Olusola O. Adetoro, Omolara Aremu, Olusegun Awotoye, Edward Missanjo, Maggie Munthali, Emmanuel F. Nzunda, Amri S. Yusuph, Pablo Miguel, Tiago Schuch Lemos Venzke, Adão Pagani Junior, José Vitor Peroba Rocha, Jeferson Diego Leidemer, Luiz Fernando Spinelli Pinto, Lizete Stumpf, Mélory Maria Fernandes de Araujo, Stefan Domingues Nachtigall, Maria Bertaso de Garcia Fernandez, Maurício Silva de Oliveira, Giovana Milech Robe, Pavel Samec, Aleš Kučera, Gabriela Tomášová, Gillian Eastwood, Manuel A. Barrios-Izás, Steven N. Winter, Sankaran Kavileveettil, Sandeep Sasidharan, Kenneth Peprah, Raymond Aabeyir, Gervase Kuuwaabong

© The Editor(s) and the Author(s) 2023

The rights of the editor(s) and the author(s) have been asserted in accordance with the Copyright, Designs and Patents Act 1988. All rights to the book as a whole are reserved by INTECHOPEN LIMITED. The book as a whole (compilation) cannot be reproduced, distributed or used for commercial or non-commercial purposes without INTECHOPEN LIMITED's written permission. Enquiries concerning the use of the book should be directed to INTECHOPEN LIMITED rights and permissions department (permissions@intechopen.com).

Violations are liable to prosecution under the governing Copyright Law.



Individual chapters of this publication are distributed under the terms of the Creative Commons Attribution 3.0 Unported License which permits commercial use, distribution and reproduction of the individual chapters, provided the original author(s) and source publication are appropriately acknowledged. If so indicated, certain images may not be included under the Creative Commons license. In such cases users will need to obtain permission from the license holder to reproduce the material. More details and guidelines concerning content reuse and adaptation can be found at <http://www.intechopen.com/copyright-policy.html>.

Notice

Statements and opinions expressed in the chapters are these of the individual contributors and not necessarily those of the editors or publisher. No responsibility is accepted for the accuracy of information contained in the published chapters. The publisher assumes no responsibility for any damage or injury to persons or property arising out of the use of any materials, instructions, methods or ideas contained in the book.

First published in London, United Kingdom, 2023 by IntechOpen
IntechOpen is the global imprint of INTECHOPEN LIMITED, registered in England and Wales, registration number: 11086078, 5 Princes Gate Court, London, SW7 2QJ, United Kingdom
Printed in Croatia

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library

Additional hard and PDF copies can be obtained from orders@intechopen.com

Forest Degradation Under Global Change

Edited by Pavel Samec

p. cm.

This title is part of the Sustainable Development Book Series, Volume 3

Topic: Climate Change and Environmental Sustainability

Series Editor: Usha Iyer-Raniga

Topic Editor: Yixin Zhang

Print ISBN 978-1-80356-794-5

Online ISBN 978-1-80356-795-2

eBook (PDF) ISBN 978-1-80356-796-9

ISSN 2753-6580

Contents

Preface	XIII
Section 1	
General Forest Degradation	1
Chapter 1	3
Perspective Chapter: Forest Degradation under Global Climate Change <i>by Sandeep Sasidharan and Sankaran Kavileveetil</i>	
Chapter 2	19
Soil Degradation Processes Linked to Long-Term Forest-Type Damage <i>by Pavel Samec, Aleš Kučera and Gabriela Tomášová</i>	
Section 2	
Impacts of Climate Change	39
Chapter 3	41
Impact of Climate Variability on Forest Vegetation Zones in Malawi <i>by Edward Missanjo and Maggie Munthali</i>	
Chapter 4	53
Forest Degradation in Tanzania: A Systematic Literature Review <i>by Emmanuel F. Nzunda and Amri S. Yusuf</i>	
Section 3	
Impacts of Human Landuse	69
Chapter 5	71
Degradation of Forest Reserves in Asunafo Forest District, Ghana <i>by Kenneth Pephrah, Raymond Aabeyir and Gervase Kuuwaabong</i>	
Chapter 6	93
Assessment of Diversity, Growth Characteristics and Aboveground Biomass of Tree Species in Selected Urban Green Areas of Osogbo, Osun State <i>by Omolara Aremu, Olusola O. Adetoro and Olusegun Awotoye</i>	

Chapter 7

107

Physical Quality of Soils in a Toposequence of a Forest Fragment under Livestock Activity in a Watershed in South Brazil

by Tiago Schuch Lemos Venzke, Pablo Miguel, Adão Pagani Junior, José Vitor Peroba Rocha, Jeferson Diego Leidemer, Stefan Domingues Nachtigall, Mélory Maria Fernandes de Araujo, Lizete Stumpf, Maria Bertaso de Garcia Fernandez, Maurício Silva de Oliveira, Giovana Milech Robe and Luiz Fernando Spinelli Pinto

Chapter 8

127

Drugs and Biodiversity Loss: Narcotrafic-Linked Landscape Change in Guatemala

by Steven N. Winter, Gillian Eastwood and Manuel A. Barrios-Izás

Preface

Human activities cause the degradation of innumerable ecosystems. In particular, forest degradation leads to far-reaching nutrient cycle changes throughout all environments. Reversible and irreversible degradation processes cause ecosystem self-regulation to decline. While reversible processes tend to be a structural simplification and a decrease in native forest biodiversity under the preservation of the ecosystem's abiotic components, irreversible processes alter the character of the potential natural vegetation due to climate change, acidification, eutrophication, and soil erosion. Subsequent secondary ecosystems already affect global change processes in an uncertain way. The book discusses the irreversible degradation processes of forests worldwide.

This book is organized into three parts on general, climatic, and land-use degradative effects. Global environmental change involves a related sequence of the biophysical, ecosystem, and socioeconomic alterations that damage the life-sustaining abilities of the planet. Current global change is caused by the human transformation of the natural environment, but also by the reaction of human communities to the induced modifications. Human transformations of the natural environment are concentrated in a critical zone. The most vulnerable part of the terrestrial critical zone is composed of soil (pedosphere), which includes interfaces among the atmosphere, hydrosphere, lithosphere, and biosphere.

The most serious soil damage is due to land-use modifications after deforestation. While mere forest felling only gives rise to reversible changes in ecosystem functions, the combination of forest felling with soil erosion or conversion for the need of subsequent management utilization generates irreversible ecosystem degradation. Deforestation is instantly followed by declining evaporation and soil loss. The imminent consequences of deforestation are gradually leading to regional climate change, the loss of ecosystem recoverability, and uninhabitable landscapes. Ultimately, landscape uninhabitability is caused by uncontrollable soil erosion because of surface exposure to wind and landslides of weathered rocks impoverished from organic binders. Thus, this book discusses characteristics of human activities that disrupt the landscape and highlights the need to restore wildlife.

Pavel Samec

Faculty of Forestry and Wood Technology,
Department of Geology and Soil Science,
Mendel University,
Brno, Czech Republic

Chapter 2

Soil Degradation Processes Linked to Long-Term Forest-Type Damage

Pavel Samec, Aleš Kučera and Gabriela Tomášová

Abstract

Forest degradation impairs ability of the whole landscape adaptation to environmental change. The impacts of forest degradation on landscape are caused by a self-organization decline. At the present time, the self-organization decline was largely due to nitrogen deposition and deforestation which exacerbated impacts of climate change. Nevertheless, forest degradation processes are either reversible or irreversible. Irreversible forest degradation begins with soil damage. In this paper, we present processes of forest soil degradation in relation to vulnerability of regulation adaptability on global environmental change. The regulatory forest capabilities were indicated through soil organic matter sequestration dynamics. We divided the degradation processes into quantitative and qualitative damages of physical or chemical soil properties. Quantitative soil degradation includes irreversible loss of an earth's body after claim, erosion or desertification, while qualitative degradation consists of predominantly reversible consequences after soil disintegration, leaching, acidification, salinization and intoxication. As a result of deforestation, the forest soil vulnerability is spreading through quantitative degradation replacing hitherto predominantly qualitative changes under continuous vegetation cover. Increasing needs to natural resources using and accompanying waste pollution destroy soil self-organization through biodiversity loss, simplification in functional links among living forms and substance losses from ecosystem. We concluded that subsequent irreversible changes in ecosystem self-organization cause a change of biome potential natural vegetation and the land usability decrease.

Keywords: global environmental change, pollution, nitrogen deposition, deforestation, soil self-organization

1. Introduction

Human activity induces degradation of many ecosystems, of which the forest degradation results in far-reaching alterations in nutrient cycles in other related types of the environment. The forest degradation, including impacts on ecosystem functions, is intensified by terrestrial environmental change. Forests are most affected by felling and critical loads of pollution. Both processes may be characterized by negative impacts on soil which subsequently causes a decline in the forest natural ability to regenerate [1]. The damage of forest soils signifies that the development of potential natural vegetation is endangered. Damaged forest soils do not allow restoration of

original plant community due to disturbed mitigation of environmental fluctuations. Soil capability to mitigate environmental fluctuations resides in uninterrupted cycles of organic matter and continuous fertility. The disruption of forest soil nutrient cycles disadvantages management utilization including sustainable landscape management [2].

Global environmental change involves a related sequence of biophysical, ecosystem and socio-economic alterations that damage life-sustaining abilities of the planet [3]. The current global change is caused by human transformation of the natural environment, but also by the reaction of human communities to the induced modifications. Human transformations of natural environment are concentrated in a critical zone. The critical zone is range among sphere interfaces on the Earth's surface, where human changes in structure, chemical composition, radiation balance and biodiversity extend [4]. The most vulnerable part of the terrestrial critical zone is composed of soil (pedosphere), which includes interfaces among atmosphere, hydrosphere, lithosphere and biosphere. For these reasons, the soil damage has been altering character of the entire ecosystem over a long period [5].

The most serious soil damage is due to land use modifications after deforestation. While mere forest felling only gives rise to reversible changes in ecosystem functions, the combination of forest felling with soil erosion or conversion for the need of subsequent management utilization generates irreversible ecosystem degradation. Deforestation is instantly followed by declining evaporation and soil loss. The imminent consequences of deforestation are gradually leading to the regional climate change, the loss of ecosystem recoverability and uninhabitable landscape [6]. Regional climate change is mainly caused by reduction of water cycle between the lower-lying areas with higher evaporation and the higher-lying areas with higher atmospheric precipitation in the catchment. The evaporation reduction after deforestation is not sufficient to create cloudiness to make surfaces cooler. Subsequent decrease in precipitation over the higher-lying parts of the river basin deepens water shortage as well as further evaporation decrease in the lower-lying parts [7]. The forest ecosystem recoverability loss is caused mainly due to depletion of the organic matter from the exposed soils, which stimulates germination of tree seeds by means of hormonal effects and moisture retention [8]. Ultimate landscape uninhabitability is caused by uncontrollable soil erosion as a result of surface exposure to wind and landslides of weathered rocks impoverished of organic binders [9].

Forest ecosystem restoration is made impossible within the recent global change, except for soil erosion by pollution. Nitrogen pollution from industry and agriculture has become a major environmental driver of the forest growth [10]. However, atmospheric pollution with the available nitrogen forms is manifested contradictorily within different soil types in forests. The forests situated on optimally fertile soils were generally favorably affected by nitrogen pollution while the predominant forests located on poor soils were damaged. On the one hand, adequate nitrogen intake supports plant growth and, on the other hand, it increases demands on other mineral resources which are declining as a result of human changes in the environment [11]. The unnaturally increasing disparity between plant demands and dwindling nutrient resources causes growth decrease and gradual ecosystem degradation even in hitherto unspoiled areas [12]. Even though the largest nitrogen deposition occurs in the vicinity of pollution sources with lower precipitation, higher concentrations of available nitrogen in wet deposition acidify ecosystems significantly. Approximately 70–80% of nitrogen released from industrial products falls back to the Earth's surface [13]. Of the nitrogen inputs, 5% penetrates the groundwater, 12%

is released into the atmosphere, 30% is immobilized in soil organic matter and 53% is removed with the crop. The utilization of nitrogen by the plant production is still declining, whereas the rate of nitrogen losses by leaching and gasification as well as immobilization in the soil increases in proportion to the amount of fertilizers [14]. Nitrogen supplied to the soil by means of fertilizers results in faster depletion of available bases, making the soil more susceptible to acidification [15].

The environmental nitrogen load is becoming an increasingly important driver of the global ecosystem change as it has exceeded the critical level in large areas of most continents [16]. Exceeding critical nitrogen loads extended plant susceptibility to drought [2, 17, 18]. The widespread plant susceptibility is compound of growing sensitivity of terrestrial ecosystems to climate change. Subsequently, the processes of the climate change and alterations in complex growth conditions for plant communities lead to a deviation in development of prospective natural vegetation or to biome alteration [19]. Therefore, the soil protection is becoming a tool to mitigate the effects of the global terrestrial change maintaining ecosystem link among forests, water cycle and human civilization [5].

2. Forest soil degradation processes

2.1 Impacts on self-organization

Soil degradation may be ranked to one of the most dangerous human activities on the Earth's surface because soil is not instantly renewable. Degradation commences with vegetation coverage damage as a result of which evaporation decreases. The evaporation diminution foreshadows regional warming which contradictorily results in an intensification of water cycle into short, intense rainfall more frequently accompanied by soil drift or even flash floods [20]. Soil degradation affects ecosystem self-organization. The disruption of soil self-organization is initiated with the decrease in the diversity of functional connections within microbial communities. Disintegration of soil functional interconnectedness involves biodiversity loss and substitution of symbioses for decomposers (saprophytes) that do not exchange available nutrients among organisms, but cause leakage of substances from the ecosystem [21]. Forest soil degradation destroys irreplaceable natural values that improve adaptability of cultural landscape to climate change [22]. The disruption of soil self-organization damages both the continuity of crop production and success of ecosystem restoration.

Soil degradation is divided into quantitative or qualitative one (**Figure 1**). Quantitative soil degradation represents the physical loss of a soil body. Qualitative soil degradation involves unfavorable alterations in soil physical or chemical properties that limit ecosystem functions [23]. Soil losses occur through claim, erosion or desertification:

- Claim is usually accompanied by soil sealing, where the land is either merely covered or removed and replaced with building materials. The claim completely destroys soil infiltration capacity. As a result of clearing, the radiation balance and heat capacity alter locally and surface runoff increases sharply [24].
- Soil erosion is surface soil drift by gravitational shifts, water or wind. The human activity has intensified soil erosion after vegetation removal, excessive grazing and inappropriate tillage, developments affecting landscape and pollution




Figure 1. Division of soil degradation processes along quantitative and qualitative impacts on physical or chemical properties.

which stopped formation of organo-mineral particles aggregating the soil into more cohesive peds (**Figure 2**). The vulnerability through erosion (erodibility) depends on weatherability of soil-forming substrate, soil cohesion, climate and land use (**Table 1**).

- Desertification is unnatural spread of wastelands after permanent vegetation removal. Causes of unnatural desertification are mainly disproportionate grazing, fires and erosion followed by loss of soil water retention capacity. Deserts spread the fastest in areas naturally adapted to seasonal drought [25]. Approximately 10–20% of the world's semi-deserts and steppes are threatened by desertification. The accompanying phenomena of desertification are decrease in groundwater levels or salinization which make it impossible to restore vegetation and lead to wasteland homeostasis [9].

Qualitative soil degradation is produced by excessive losses of the organic matter, the reduction of the biological activity, acidification, contamination, technological compaction (pedocompaction), technical or wind salinization and



Figure 2. Coupled occurrences of soil erosion and spreading of desert in arid environments seriously threat forest restoration due to water availability decrease.

Erodibility	Parent rock	Soil units
Very easy	Eolian deposits	Leptosols, Retisols, Anthroposols
Easy	Clayey shales, basaltic tuffs	Leptosols, Luvisols, Cambisols
Medium	Carbonate deposits, sandstones	Calcaric Leptosol, Chromic Cambisol, Ferralic Podzol
Medium hard	Breccias, graywackes, phyllites, andesites	Stagnic Podzol, Stagnosols, Stagnic Cambisol
Hard	Metamorphites, basalts, diorites	Gleysols, Stagnic Luvisol, Haplic Cambisol
Very hard	Igneous rocks, shales	Epi-humic Cambisol, Vertic Chernozem, Histosols

Table 1. The vulnerability of forest soils by erosion (erodibility) along various parent rocks and developed soil units.

technical modifications of soil properties. Although qualitative soil degradation potentially occurs in much smaller areas than quantitative one and its effects are usually reversible, they have a similar overlap on landscape functions in the form of reduced water retention capacity and biodiversity, increased runoff and substance imbalances.

Degradation of forest soils is distinctive mainly to the qualitative damage. Qualitative degradation in forests prevails due to the long-term growth of continuous tree species communities. Tree species instantly impede quantitative damage to soils; on the other hand, periodic windthrows during storms mingle the mass among soil horizons, as well as move the soil down the slope. These post-disturbance movements




Figure 3. A series of differently degraded soil bodies in mountain conditions: introskeletal erosion in Dystric Hyperskeletal Leptosol (A); surface scarification of Entic Podzol (B) and accumulated Spolic Garbic Urbic Technosol (C).

of earth bodies divide microrelief and homogenize soil properties, but at the same time their arrangement is concentrated along the effects of individual global climate changes [26]. However, predominant qualitative degradation of forest soils is manifested by deterioration of physical or chemical properties of solid soil bodies due to external human activities (Figure 3) [27].

Deterioration of forest renewal as a consequence of qualitative soil degradation commences with fertility change. Forest felling is accompanied by accelerated leaching of nitrogen substances which can only be ceased by sufficient available calcium [28]. Leaching is preceded by increase in C/N which indicates decrease in ability of soil organic matter to bind mineral nutrients. Soils damaged by compaction of the profile middle part, texturally significantly differentiated or hydromorphic, are exposed to a slow-motion water flow, which expels air for the root growth and similarly increases C/N [29]. The root systems grow merely shallowly with water stagnation and the nutrient loss, making forest stands more susceptible to soil moisture fluctuations [30]. Thus, felling of forests threatened by qualitative erosion impairs the ecosystem ability to restore as a result of exposure to episodic drought [2].

2.2 Physical degradation

Degradation of soil physical properties includes structural damage, pore loss and compaction. The processes of physical soil damage result in both loss of water retention capacity and humus loss by introskeletal erosion. The decline in soil water retention capacity is usually caused by repeated heavy machinery moving. Heavy machinery moving worsens soil aeration and water permeability. Reduced aeration is reflected in the decrease of blank spaces for plant roots and the consequent reduction in the biological activity.

Forest soils are less endangered by physical degradation than agricultural ones owing to dampening effect of surface humus. Nevertheless, topsoil compaction can initiate introskeletal erosion. The mechanical degradation threat is descending from Histosols and gleyed soils to granularly light drying soil bodies [31]. The risks by mechanical damage to the forest soils vary along the grain-size composition, relief exposure and groundwater level (Table 2) [32].

Risk	Soil series	Inprint depth	Consistency	Critical pressure (kPa)	
				Dry conditions	Wet conditions
Very high	Histosols+Gleysols	≥35	Cohesionless	30–50	5–12
High	Stagnosols+stagnic soil groups	25–35	Viscous	50–140	12–22
Medium	Stagnic Cambisols-Stagnic Luvisols-Fluvisols	15–25	Crombly	140–300	18–50
Moderate	unhydromorphic Cambisols+Luvisols+ Chernozems+ Regosols	7–15	Cohesive	300–600	50–80
Insignificant	Leptosols+Podzols	<7	Skeletal	600	80–120

Table 2.
 Characteristics of forest soil compaction risk after logging machinery movement.

- A very high risk results from high groundwater level and incoherent soil-forming substrate. Forests endangered by a very high risk of soil erosion are mostly found on Histosols or Gleysols, but also on water-affected Luvisols, Arenosols or Podzols.
- A high risk emerges from extremely developed hydromorphic features related to heavy skeletalness of soil bodies. Forests exposed to high erosion risk are covering Planosols, Stagnosols and Gleysols, including gleyed subtypes on clayey shales or claystones the most.
- A medium risk is associated with the medium level of forest site gleyfication. Forests at the medium soil erosion risk are located on Stagnic Cambisols, Stagnic Luvisols or Stagnic Fluvisols.
- A moderate risk is related to site desiccation. Forests moderately endangered by erosion means may be found on Cambisols, Luvisols, Chernosols or on Fluvisols developed from sandy substrates.
- An insignificant risk is conditioned by soil cohesion, medium skeletalness and merely slightly by sloping relief. Forests insignificantly exposed to soil erosion occur at unexposed sites constituted by Leptosols, Cambisols, Podzols or by Chernosols.

Introskeletal erosion represents a predominantly vertical subsidence of fine-grained soil particles through blank spaces among skeleton to the base of rock mantle. The introskeletal erosion risk resides in unstable occurrence of surface humus. Introskeletal erosion is triggered after removal of the vegetation cover in the exposed sites. Its result is the loss of whole fine-grained matter, followed by impossibility of restoring plant community and permanent exposure of relief [33]. The

threat to the site by introskeletal erosion is distributed along exposure of relief and soil skeletability [34]:

- An extreme risk accompanies periglacial brash in arcto-alpine conditions. Extremely endangered sites are only merely sparsely populated by forests. Emerging plant communities are very sensitive to any changes in growth conditions, so they require consistent protection.
- A very strong risk accompanies shallow soils. Forests endangered by very strong risk of introskeletal erosion are most frequently found along upper tree vegetation limit that is sensitive to global warming [35].
- A strong risk is accompanied with brash or stone fields (**Figure 4**). Forests exposed to the strong risk of introskeletal erosion are mostly concentrated on long rocky slopes below the upper limit of tree species vegetation.
- A medium risk is characteristic of islet occurrences of brash on rocky slopes. Forests exposed to the medium risk of introskeletal erosion typically occur in the middle parts of mountain ranges.
- A low risk is specific for sparse outcrops of subsoil decay on medium rocky slopes. Forests threatened by erosion on a small scale occur on gentle slopes with deeply developed soils

2.3 Chemical degradation

2.3.1 Acidification

Degradation of soil chemical properties is the intensification of naturally processed weathering and substance leaching. Chemical soil degradation includes acidification, salinization and intoxication. Acidification is the most extensive process of forest soil degradation causing decline in site fertility [36]. Soil acidification is gradual decreased in neutralizing capacity. In nature, acidification is elicited mainly by water autoprotolysis, naturally acid atmospheric precipitation, organic acids activities, but also by formation of strong acids after reactions of water with atmospheric gases (CO_2 , SO_2) or with some rock-forming minerals (chlorides, sulphates or carbonates). The resulting acids (formal HCl , H_2CO_3 and H_2SO_3) can cause very intensive decomposition of original minerals into salts [37].

The intensification of soil acidification was caused by fertilization, crop cultivation and industrial pollution. Industrially emitted CO_2 , SO_2 and NO_x create formal acids and soil bases are excessively depleted to neutralize them. The base loss slows down humus formation; on the other hand, raw humus is a significant source of organic acids. The slow-motion formation of humus is reflected in decrease of organo-mineral colloid genesis as a result of which the number of binding sites for exchange cations on the active surfaces of soil particles decreases. The decomposition of variable organo-mineral colloids limits base cations exchanges to stable mineral colloids. Nonetheless, mineral colloids can capture only 0.2–25% of exchangeable cations, unlike organic particles [38].

The soil resists acidification impacts by exchange reactions between inputs of acid-forming H_3O^+ and available sources of releasable cations. Soil cation sources are



Figure 4. *Intraskelatal erosion leaves rocky flows without surface humus instead soil, where plants can to root hardly, thus forest site gets features of disperse platforms with dwarf vegetation.*

active depending on pH (**Figure 5**). Intensified acidification of forest soils is naturally slowed down either by deciduous tree species or by weathering of the soil-forming substrates. The influence of tree species predominates in surface soil horizons while the influences of soil-forming substrate predominate in the subsurface horizons. Significant acidification in surface horizons of forest soils most often affects the transitional ecosystem types [39]. Even though the mitigating effect of tree species does not overcome impacts of weathering, the optimal tree species composition actively




Figure 5. *Intervals of soil acidity (pH) and organic matter C/N ratio divide trophic (A – Oligotrophic; AB – Oligomesotrophic; B – Mesotrophic; BC – Mesotrophically nitrophilous; C – Nitrophilous; CD – Nitrophilous-base; BD – Mesotrophically base; D – Base) series among zones buffering acidification through specific neutralization. Data according to [39].*

reducing C/N prolongs weathering effects. On the other hand, soil cation release by weathering maintains intensified acidification as a reversible process. Weathering counteracts acidification by means of electrochemically controlled soil-forming substrate decomposition [12, 40, 41]:

1. **The carbonate zone** (pH > 6.8) is employed by dissolving the CO_3^{2-} compounds, whereby the incoming H^+ is neutralized into soluble salts. The consequence of these acid-base reactions is a gradual loss of carbonates by dissolution and leaching, which can only be forestalled in soils formed directly from carbonate substrates and rocks.
2. **The silicate zone** (pH 5.0–6.8) occurs either within soils from which carbonates have already been leached or where silicates predominate. Acids cause decomposition of silicates from which base cations are released and deuterogenous clay minerals are formed.
3. **The exchange zone** (pH 4.2–5.0) may be found in those soils where there is a disproportion between base cations released during weathering of silicates and H^+ inputs. The excess of entering H^+ is trapped on surface of organic colloids to release bases.
4. **The aluminum zone** (pH 3.0–4.2) subdues effects of acidic inputs by releasing Al^{3+} in the presence of sesquioxides with simultaneous formation of organic complexes. Soil fertility decreases, nutrients are further leached and the biological activity decreases.
5. **The iron zone** (pH < 3.0) occurs in those soils where acidic inputs are subdued by the dissolution of iron oxides, Fe^{3+} migration and destruction of clayey minerals. Nutrients are excessively leached out from these soil bodies, the concentration of toxic substances in soil profile increases and the biological activity is usually concentrated only into raw surface humus.

The unnatural decomposition of soil minerals triggers irreversible acidification. Acidification may be mitigated merely after the removal of acidifying substances sources. The acidification of forest soils affected exchange zone the most, switching to active aluminum zone [42]. The damage to the forest ecosystem by release of active Al^{3+} followed due to occupation of exchange sites on soil particle surfaces instead of bases, the lack of which limited root growth. The roots were concentrated shallowly below the surface so that new focal points of biological activity and humus ceased to form deeper in the soil [43]. Introducing the other side of the fact, the marginally widespread transition from the aluminum zone to iron one was ensued by loss at the ability of forest ecosystems to restore from the damage (Figure 6).

Air pollution has significantly accelerated soil acidification, especially in the areas of forests transformed into homogeneous stands of coniferous tree species. While cultivation of homogeneous coniferous forests homogenized formation of acidic humus causing micropodzolization and increased base cation leaching, the pollution after acid deposition reduced not only the forest increment but also decomposition of organic matter [44]. Forest increment was reduced by direct damage to the assimilation apparatus, by stimulating sensitivity to seasonal drought or frost and by reduction in soil symbioses mediating nutrient deficiencies. The decline of mycorrhizal



Figure 6. Irreversible damaged forests are characteristic by predominantly dead tree storey and by absent young woods due to lost soil organic matter irreplaceably stabilizing moisture during seed germination.

fungi was followed by increase in frequency of saprophytic to saproparasitic fungi, which diverted organic matter decomposition to complete leaching from the ecosystem [45]. The susceptibility of mycorrhizal symbioses to pollution resulted in limited accessibility to phosphorus necessary for nucleic acid synthesis [46]. The disturbed phosphorus cycle triggered decrease in increment as well as seed germination leading to forest self-organization loss [47].

2.3.2 Salinization

Soil salinization is the process of accumulating surpluses of mineral salts. Salinization of forest soils is a rare phenomenon, but it threatens 23% of agricultural land, mainly in arid areas [48]. Forest soils are salinized in areal or linear extent. Areal salinization is caused by high groundwater mineral levels, the use of saline water for irrigation, waste materials for fertilization or deposition of solids. Linear salinization occurs alongside roadsides maintained by chemical salting during winter or along river banks. The recent climate change is expanding areas of salinized soils with rising sea level along the coast or estuaries. On the other hand, the natural risk of soil acidification subdues consequences of salinization [49].

The impacts of salinization in forests are associated with extreme soil chemical properties. Salinization highlights malfunctions of water and nutrient uptake by plants. Above all, the disproportionate sodium input (sodification) disrupts rations among exchangeable bases in the soil environment, thereby disrupting effects of alkalization on soil structure. Significant Na^+ inputs displace other cations from soil sorption complex and disperse soil particles. Sodium displacement of cations results in deficient nutrition, but at the same time crushes soil structure, thus water availability fluctuates. Sodium surplus in plant tissues reduces osmotic pressure, whereby cells lose ability to absorb other substances from soil solution [27]. While conifers are

susceptible to soil salinization, deciduous tree species are tolerant to it. The younger plants are more susceptible than the older ones.

Areal forest salinization is most at risk in floodplains due to variability of water flow. The regulation of water flow caused groundwater level fall in some river basins while it resulted to water level permanent increase in some other ones. The groundwater level decline was typically ensued by ecosystem desiccation due to the fact that riparian forests are mostly located in submontane locations with insufficient precipitation [50]. By contrast, rising groundwater levels after water regulation meant change in availability of mineral ions, with impacts on soil microbial activity and ability of the ecosystem to sequester carbon. The increase in level of saline water inflicts decrease in soil microbial activity and consequently decrease in vegetation growth [51]. On the one hand, decreases in growth processes are caused by loss of oxygen in soil environment, on the other hand, by increasing concentrations of Na^+ , Cl^- , SO_4^{2-} ions, including Fe and Mn compounds. In particular, SO_4^{2-} in the soil solution is converted to toxic sulphide when there is a lack of oxygen. Although Fe and Mn are biogenic elements that catalyze soil organic matter decomposition, bound in sulphides block microbial metabolism [52]. The main forest salinization danger with groundwater is inability to adapt on climate change due to spread of microaerobic conditions [53, 54].

2.3.3 Intoxication

Intoxication of soils with heavy metals, radioactive or petroleum substances is a rare but very hazardous process of cumulative pollution. Especially, heavy metals merely slowly participate in biogeochemical cycles and accumulate in the ecosystem because they are either microbiogenic (Cu and Zn), or xenobiotic (e.g. Cd, Co, Pb, Hg, Ni).

Soil intoxication occurs by deposition means. Sources of cumulative pollution are point or dispersed ones. The point sources of heavy metals are smelters, thermal power stations or municipalities by watercourses. The dispersed sources are represented by polluted water, inappropriate distribution of industrial or sewage sludge or operation of internal combustion engines. Fluvisols, which are among the most intoxicated soils, are usually located between watercourses and agricultural soils [48].

Xenobiotics are toxic to most organisms. They mainly affect energy balance of living cells and their division. Heavy metals mainly bring about halting respiration as a consequence of interactions with SH- groups at intracellular enzymes and their complexes, they disrupt semipermeability of cell membranes and their proton gradient. Soil environment pollution with heavy metals significantly reduces density of microbial occurrences and directly damages plants. The rate of soil contamination damages microbial activity significantly more than the differences in heavy metal contents among different sites [55]. However, the decline of susceptible species is being replaced by expansion of resistant species populations, including pests abundantly infesting damaged plants [56].

Resistant phytophagous arthropods adapt to the environment contaminated with heavy metals by searching for less contaminated nourishment, sufficient release of metals in excrements, by sloughing or by means of other tissues (in the adipose body, epithelium of the digestive tract) [57]. The importance of phytophagous insects for the movement of heavy metals in ecosystem lies in the fact that these invertebrates are an important link in food web that receive toxic substances directly from plants, especially Cd and Zn [58, 59], but no Ni and Fe [60].

The risk of heavy metal accumulation with toxic manifestations threatens not only tree species, but also predators. Numerous ground beetles (Carabidae) and ants (Formicidae) are mainly food-bound to phytophagous insects, earthworms, countless larvae and springtails [61]. On the other hand, the immobilization of heavy metals takes place in cells of specialized metallogenic microorganisms by binding to metallothionein-based amino acids. Immobilization of heavy metals in amino acids changes course of humus formation. The rate of biosorption on the soil microbial active surfaces typically decreases in order $Zn > Cd > Pb > Cu > Cr$ [62]. Alterations in the forest ecosystems as a result of heavy metal pollution include (**Figure 7**):

- the reduction of phospholipid fatty acids content in the bacterial cells and raw humus. Even though the total content of phospholipid acids is directly proportional to ATP synthesis, it decreases under the toxic load. The consequence of these processes is alterations in overall functional diversity of microbial community and inability to decompose deposition by aromatic hydrocarbons [63].
- promoting release of mobile humus substances and decrease of insoluble substances. On the other hand, the great ability of Pb to form complexes with insoluble humus substances maintains its immobilization while Cd and Zn tend to form soluble complexes. Although more heavy metals are bound to soluble humus substances than to insoluble organic compounds, the greater release of soluble compounds also contributes to pollutant migration in the soil and to their penetration into groundwater [64].

Petroleum products belong to secondary persistent organic compounds, similar to benzo(a)pyrene, polychlorinated dibenzo-p-dioxins or dibenzofurans, which are removed from the soil for more than 2 years [65]. Like benzo(a)pyrene, they consist of polycyclic aromatic hydrocarbons that directly harm the health of organisms. Oil pollution is significantly more caused by human activity than by natural (geogenic) sources. It begins with mining, combustion of petroleum products (fossil fuels), accidental or operational spills and corrosion of industrial materials. Oil products in the forest ecosystem load surface humus the most, which at the same time prevents their



Figure 7. Forests dying at regions loaded by acid deposition were transformed to substitute stands of resistant introduced tree species which have provided cover for regeneration of indigenous forest communities after pollution decrease.

penetration into deeper occurring soil. The load of surface humus decreases activity of soil microorganisms. The decrease of soil biological activity is mostly caused by aromatic nuclei imitating lignin, which either block formation of amino acids or replace carbon compounds in fungi [66]. Subsequently, the humus decomposition is disrupted, foreshadowing disruption of processes to get available nutrients from the soil. Nevertheless, the load of petroleum substances is irregularly concentrated in surroundings of industrial areas and vertically along different intensities of wood logging in floodplains, hilly countries, highlands and high-mountain forests [67].

Acknowledgements

The study was supported by projects LM2018123 CzeCOS of the Ministry of Education, Youth and Sports of the Czech Republic and 952314 ASFORCLIC of the European Union's Horizon 2020 Programme for Research & Innovation.

Author details


Pavel Samec^{1,2*}, Aleš Kučera¹ and Gabriela Tomášová¹

1 Faculty of Forestry and Wood Technology, Department of Geology and Soil Science, Mendel University in Brno, Brno, Czech Republic

2 Global Change Research Institute CAS, Brno, Czech Republic

*Address all correspondence to: pavel.samec@mendelu.cz

IntechOpen

© 2022 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, Tyukavina A, et al. High-resolution global maps of 21st-century forest cover change. *Science*. 2013;**342**:850-853. DOI: 10.1126/science.1244693
- [2] Bernal S, Hedin LO, Likens GE, Gerber S, Buso DC. Complex response of the forest nitrogen cycle to climate change. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;**109**:3406-3411. DOI: 10.1073/pnas.1121448109
- [3] Brovkin V, Brook E, Williams JW, Bathiany S, Lenton TM, Barton M, et al. Past abrupt changes, tipping points and cascading impacts in the earth system. *Nature Geoscience*. 2021;**14**:550-558. DOI: 10.1038/s41561-021-00790-5
- [4] Richter DB, Mobley ML. Monitoring Earth's critical zone. *Science*. 2009;**326**:1067-1068. DOI: 10.1126/science.1179117
- [5] Kučera A, Samec P, Bajer A, Skene KR, Vichita T, Vranová V, et al. Forest soil water in landscape context. In: Datta R, Meena RS, editors. *Soil Moisture Importance*. London, UK: InTechOpen; 2021
- [6] Woo SW. Forest decline of the world: A linkage with air pollution and global warming. *Africal Journal of Biotechnology*. 2009;**8**:7409-7414
- [7] Lawton RO, Nair US, Pielke RA Sr, Welch RM. Climatic impact of tropical lowland deforestation on nearby montane cloud forests. *Science*. 2001;**294**:584-587. DOI: 10.1126/science.1062459
- [8] Lucier A, Ayres M, Karnosky D, Thompson I, Loehle C, Percy K, et al. Forest responses and vulnerabilities to recent climate change. In: Seppälä R, Buck A, Katila P, editors. *Adaptation of Forests and People to Climate Change – A Global Assessment Report*. Vienna, Helsinki: IUFRO; 2009. pp. 29-52
- [9] Margono BA, Potapov PV, Turubanova S, Stolle F, Hansen MC. Primary forest cover loss in Indonesia over 2000-2012. *Nature Climate Change*. 2014;**4**:730-735. DOI: 10.22146/ijg.12496
- [10] Etzold S, Ferretti M, Reinds GJ, Solberg S, Gessler A, Waldner P, et al. Nitrogen deposition is the most important environmental driver of growth of pure, even-aged and managed European forests. *Forest Ecology and Management*. 2020;**458**:117762. DOI: 10.1016/j.foreco.2019.117762
- [11] Nosengo N. Fertilized to death. *Nature*. 2003;**425**:894-895. DOI: 10.1038/425894a
- [12] Schröder W, Nickel S, Jenssen M, Riediger J. Methodology to assess and map the potential development of forest ecosystems exposed to climate change and atmospheric nitrogen deposition: A pilot study in Germany. *Science of the Total Environment*. 2015;**521**:108-122. DOI: 10.1016/j.scitotenv.2015.03.048
- [13] Galloway JN, Schlesinger WH, Hiran Levy II, Michaels A, Schnoor JL. Nitrogen fixation: Anthropogenic enhancement – Environmental response. *Global Biogeochemical Cycles*. 1995;**9**:235-252. DOI: 10.1029/95GB00158
- [14] Goulding KWT, Bailey NJ, Bradbury NJ, Hargreaves P, Howe M, Murphy DV, et al. Nitrogen deposition and its contribution to nitrogen cycling and associated soil processes. *The New*

- Phytologist. 1998;**139**:49-58.
DOI: 10.1046/j.1469-8137.1998.00182.x
- [15] Oulehle F, Evans CD, Hofmeister J, Krejčí R, Tahovská K, Persson T, et al. Major changes in forest carbon and nitrogen cycling caused by declining sulphur deposition. *Global Change Biology*. 2011;**17**:3115-3129.
DOI: 10.1111/j.1365-2486.2011.02468.x
- [16] Gruber N, Galloway JN. An earth-system perspective of the global nitrogen cycle. *Nature*. 2008;**451**:293-296.
DOI: 10.1038/nature06592
- [17] Westling O, Fagerli H, Hellsten S, Knulst JC, Simpson D. Comparison of modelled and monitored deposition fluxes of Sulphur and nitrogen to ICP – Forest sites in Europe. *Biogeosciences Discussions*. 2005;**2**:933-975. DOI: 10.5194/bg-3-337-2006
- [18] Vacek S, Vacek Z, Remeš J, Bílek L, Hůnová I, Bulušek D, et al. Sensitivity of unmanaged relict pine forest in the Czech Republic to climate change and air pollution. *Trees*. 2017;**31**:1599-1617. DOI: 10.1007/s00468-017-1572-0
- [19] Parmesan C, Yohe G. A globally coherent fingerprint of climate change impacts across natural systems. *Nature*. 2003;**421**:37-42. DOI: 10.1038/nature01286
- [20] Blöschl G, Kiss A, Viglione A, Barriendos M, Böhm O, Brázdil R, et al. Current European flood-rich period exceptional compared with past 500 years. *Nature*. 2020;**583**:560-566.
DOI: 10.1038/s41586-020-2478-3
- [21] Abbasian F, Lockington R, Megharaj M, Naidu R. The biodiversity changes in the microbial population of soils contaminated with crude oil. *Current Microbiology*. 2016;**72**:663-670.
DOI: 10.1007/s00284-016-1001-4
- [22] Bergstrom DM, Wienecke BC, van den Hoff J, Hughes L, Lindenmayer DB, Ainsworth TD, et al. Combating ecosystem collapse from the tropics to the Antarctic. *Global Change Biology*. 2021;**27**:1692-1703. DOI: 10.1111/gcb.15539
- [23] Lal R. Soil quality and sustainability. In: Lal R, Blum WH, Valentine C, Steward BA, editors. *Methods for Assessment of Soil Degradation*. Boca Raton: CRC Press; 1998. pp. 547-554
- [24] Bronstert A, Niehoff D, Bürger G. Effects of climate and land-use change on storm runoff generation: Present knowledge and modelling capabilities. *Hydrological Processes*. 2002;**16**:509-529.
DOI: 10.1002/hyp.326
- [25] Trnka M, Hayes M, Jurečka F, Bartošová L, Anderson M, Brázdil R, et al. Priority questions in multidisciplinary drought research. *Climate Research*. 2018;**75**:241-260. DOI: 10.3354/cr01509
- [26] Šilhán K, Pánek T, Dušek R, Havlů D, Brázdil R, Kašičková L, et al. The dating of bedrock landslide reactivations using dendrogeomorphic techniques: The Mazák landslide, outer Western Carpathians (Czech Republic). *Catena*. 2013;**104**:1-13. DOI: 10.1016/j.catena.2012.12.010
- [27] Vavříček D, Kučera A. *Základy lesnického půdoznalství a výživy lesních dřevin*. Kostelec nad Černými lesy: Lesnická práce; 2017. p. 364
- [28] Green MB, Bailey AS, Bailey SW, Battles JJ, Campbell JL, Driscoll CT, et al. Decreased water flowing from a forest amended with calcium silicate. *Proceedings of the National Academy of Sciences of the United States of America*. 2013;**110**:5999-6003. DOI: 10.1073/pnas.1302445110

- [29] Likens GE, Bormann FH. Biogeochemistry of a Forested Ecosystem. New York: Springer-Verlag; 1995
- [30] Knapp AK, Beier C, Briske DD, Classen AT, Luo Y, Reichstein M, et al. Consequences of more extreme precipitation regimes for terrestrial ecosystems. *Bioscience*. 2008;**58**:811-821. DOI: 10.1641/B580908
- [31] Vavříček D, Ulrich R, Kučera A. Ochrana půdy v těžebně dopravní činnosti. Mendelova univerzita v Brně. 2014
- [32] Simanov V, Macků J, Popelka J. Nový návrh terénní klasifikace a technologické typizace. *Lesnictví-Forestry*. 1993;**39**:422-428
- [33] Šach F, Černohous V. Introskeletová eroze. In: Borůvka L, editor. *Pedologické dny*. 2002. pp. 129-132
- [34] Souček J, Kriegel H, Nárovec M, Šach F. Obnova lesa na lokalitách ohrožených introskeletovou erozí. *Lesnický průvodce*. 2010;**2**:1-35
- [35] Wang Z, Lyu L, Liu W, Liang H, Huang J, Zhang QB. Topographic patterns of forest decline as detected from tree rings and NDVI. *Catena*. 2021;**198**:105011. DOI: 10.1016/j.catena.2020.105011
- [36] Cosby BJ, Ferrier RC, Jenkins A, Wright RF. Modelling the effects of acid deposition: Refinements, adjustments and inclusion of nitrogen dynamics in the MAGIC model. *Hydrology and Earth System Sciences*. 2001;**5**:499-517. DOI: 10.5194/hess-5-499-2001
- [37] Bolan NS, Curtin D, Adriano DC. Acidity. In: Hillel D, editor. *Encyclopedia of Soils in the Environment*. New York: Academic Press; 2005. pp. 11-17
- [38] Ulrich B. The history and possible cause of forest decline in Central Europe, with particular attention to the German situation. In: EC, UN/ECE, *Forest Soil Condition in Europe*. Brussel, Geneva: Large-Scale Soil Survey; 1995
- [39] Samec P. Indication of forest soil fertility deviations by differences between trophic series and soil buffering: Geobiocoenological approach. *Phytopedon (Bratislava)*. 2016;**15**:1-11
- [40] Fenn ME, Huntington TG, McLaughlin SB, Eagar C, Gomez A, Cook RB. Status of soil acidification in North America. *Journal of Forest Science*. 2006;**52**(Special Issue):3-13
- [41] Ross DS, Matschonat G, Skjellberg U. Cation exchange in forest soils: The need for a new perspective. *European Journal of Soil Science*. 2008;**59**:1141-1159. DOI: 10.1111/j.1365-2389.2008.01069.x
- [42] Paoletti E, Schaub M, Matyssek R, Wissler G, Augustaitis A, Bastrup-Birk AM, et al. Advances of air pollution science: From forest decline to multiple-stress effects on forest ecosystem services. *Environmental Pollution*. 2010;**158**:1986-1989. DOI: 10.1016/j.envpol.2009.11.023
- [43] Heim A, Luster J, Brunner I, Frey B, Frossard E. Effects of aluminium treatment on Norway spruce roots: Aluminium binding forms, element distribution, and release of organic substances. *Plant and Soil*. 1999;**216**:103-116
- [44] Klimo E, Materna J, Lochman V, Kulhavý J. Forest soil acidification in the Czech Republic. *Journal of Forest Science*. 2006;**52**(Special Issue):14-22
- [45] Hagerberg D, Thelin G, Wallander H. The production of ectomycorrhizal mycelium in forests: Relation between

- forest nutrient status and local mineral sources. *Plant and Soil*. 2003;252:279-290. DOI: 10.1023/A:1024719607740
- [46] Rejšek K. Acid phosphomonoesterase activity of ectomycorrhizal roots in Norway spruce pure stands exposed to pollution. *Soil Biology and Biochemistry*. 1991;23:667-671. DOI: 10.1016/0038-0717(91)90081-T
- [47] Talkner U, Meiwes KJ, Potočić N, Seletković I, Cools N, De Vos B, et al. Phosphorus nutrition of beech (*Fagus sylvatica* L.) is decreasing in Europe. *Annals of Forest Science*. 2015;72:919-928. DOI: 10.1007/s13595-015-0459-8
- [48] FAO-ITPS. Status of the World's soil resources (SWSR) – Main report. Rome: Food and agriculture Organization of the United Nations. Intergovernmental Technical Panel on Soils. 2015
- [49] Shi WM, Yao J, Yang F. Vegetable cultivation under greenhouse conditions leads to rapid accumulation of nutrients, acidification and salinity of soils and groundwater contamination in south-eastern China. *Nutrient Cycling in Agroecosystems*. 2009;83:73-84. DOI: 10.1007/s10705-008-9201-3
- [50] Klimo E, Hager H, Matic S, Anić I, Kulhavý J. Floodplain forests of the temperate zone of Europe. Kostelec nad Černými lesy: Lesnická práce; 2008
- [51] Machado RMA, Serralheiro RP. Soil salinity: Effect on vegetable crop growth. Management practices to prevent and mitigate soil salinization. *Horticulturae*. 2017;3:30. DOI: 10.3390/horticulturae3020030
- [52] Lamers LP, Dolle GE-T, Berg STVD, van Delft SP, Roelofs JG. Differential responses of freshwater wetland soils to sulphate pollution. *Biogeochemistry*. 2001;55:87-101. DOI: 10.1023/A:1010629319168
- [53] Herbert ER, Boon P, Burgin AJ, Neubauer SC, Franklin RB, Ardón M, et al. A global perspective on wetland salinization: Ecological consequences of a growing threat to freshwater wetlands. *Ecosphere*. 2015;6:206. DOI: 10.1890/ES14-00534.1
- [54] Pechanec V, Machar I, Kiliánová H, Vyvlečka P, Seják J, Pokorný J, et al. Ranking the key Forest habitats in ecosystem function provision: Case study from Morava River basin. *Forests*. 2021;12:138. DOI: 10.3390/f12020138
- [55] Friedlová M. The influence of heavy metals on soil biological and chemical properties. *Soil & Water Research*. 2010;5:21-27. DOI: 10.17221/11/2009-SWR
- [56] Führer E. Air pollution and the incidence of forest insect problems. *Zeitschrift für Angewandte Entomologie*. 1985;99:371-377. DOI: 10.1111/j.1439-0418.1985.tb02000.x
- [57] Lindqvist L. Accumulation of cadmium, copper, and zinc in five species of phytophagous insects (Swe.). *Environmental Entomology*. 1992;21:160-163. DOI: 10.1093/ee/21.1.160
- [58] Roberts RD, Johnson MS. Dispersal of heavy metals from abandoned workings after their transference through terrestrial food chains. *Environmental Pollution*. 1978;16:293-310. DOI: 10.1016/0013-9327(78)90080-0
- [59] Nuorteva P. The Effect of Pollution in Development of Forest Insect Outbreaks. Petrozavodsk: Barents Euro-Arctic Region; 1997. pp. 248-249
- [60] Beyer WN, Pattee OH, Sileo L, Hoffman DJ, Mulhern BM.

Metal contamination in wildlife living near two zinc smelters. *Environmental Pollution*. 1985;38:63-86. DOI: 10.1016/0143-1471(85)90094-7

[61] Kula E, Purchart L. The ground beetles (Coleoptera: Carabidae) of forest altitudinal zones of the eastern part of the Krušné hory Mts. *Journal of Forest Science*. 2004;50:456-463. DOI: 10.17221/4641-JFS

[62] Muter O, Lubinya I, Millers D, Grigorjeva L, Ventinya E, Rapoport A. Cr (IV) sorption by intact and dehydrated *Candida utilis* cells in the presence of other metals. *Process Biochemistry*. 2002;38:123-131. DOI: 10.1016/S0032-9592(02)00065-1

[63] Laczko E, Rudaz A, Aragno M. Diversity of anthropogenically influenced or disturbed soil microbial communities. In: Insam H, Rangger A, editors. *Microbial Communities. Functional Versus Structural Approaches*. Berlin – Heidelberg: Springer Verlag; 1997. pp. 57-67

[64] Borůvka L, Drábek O. Podíl těžkých kovů ve frakcích huminových látek kontaminované půdy. In: Borůvka L, editor. *Pedologické dny 2002 – Degradace půdy*. ČZU v Praze; 2002. pp. 55-58

[65] Němeček J, Vácha R, Podlešáková E. Hodnocení kontaminace půd v ČR. Praha: Výzkumný ústav meliorací a ochrany půdy; 2010. p. 148

[66] Vácha R, Čechmánková J, Havelková M, Horváthová V, Skála J. Přestup polycyklických aromatických uhlovodíků z půdy do vybraných rostlin. *Chemické Listy*. 2008;11:1003-1010

[67] Borůvka L, Sáníka M, Šrámek V, Vácha R, Čechmánková J, Čupr P, et al. Methods for the Forest soils pollution assessment. *Lesnický průvodce*. 2015;12:1-63

[82] Magsalay P, Brooks T, Dutson G, Timmins R. Extinction and conservation on Cebu. *Nature*. 1995;**373**:294-294

[83] Brooks T, Balmford A. Atlantic forest extinctions. *Nature*. 1996;**380**:115-115

[84] Grelle CEDV, Fonseca GAB, Fonseca MT, Costa LP. The question of scale in threat analysis: A case study with Brazilian mammals. *Animal Conservation*. 1999;**2**:149-152

[85] Wich SA, Singleton I, Nowak MG, Utami Atmoko SS, Nisam G, Arif SM, et al. Land-cover changes predict steep declines for the Sumatran orangutan (*Pongo abelii*). *Science Advances*. 2016;**2**(3):e1500789. Available from: <https://insightcrime.org/news/brief/us-extradition-request-depicts-deep-ties-guatemala-ex-vp-zetas/>

[86] Estavillo C, Pardini R, Rocha PLBD. Forest loss and the biodiversity threshold: An evaluation considering species habitat requirements and the use of matrix habitats. *PLoS One*. 2013;**8**:e82369

[87] Ciccarone D. Heroin in brown, black and white: Structural factors and medical consequences in the US heroin market. *International Journal of Drug Policy*. 2009;**20**:277-282

[88] United Nations Office of Drugs and Crime. *World Drug Report*. United Nations; 2020

Edited by Pavel Samec

Human activities cause the degradation of innumerable ecosystems, including forests. The degradation of forests leads to far-reaching nutrient cycle changes throughout all environments. Reversible or irreversible degradation processes cause declines in an ecosystem's ability to self-regulate. Reversible processes decrease native forest biodiversity while preserving the ecosystem's abiotic components, whereas irreversible processes alter the character of the potential natural vegetation. This book discusses forest degradation in three sections on general, climatic, and land-use degradative effects.

*Usha Iyer-Raniga,
Sustainable Development Series Editor*

Published in London, UK

© 2023 IntechOpen
© Andre2013 / iStock

IntechOpen

ISSN 2753-6580

ISBN 978-1-80356-794-5



9 781803 567945