Mendel University in Brno Czech Society of Landscape Engineers – ČSSI, z.s.

Public recreation and landscape protection – with environment hand in hand?



Proceedings of the 14th Conference

Editor: Jitka Fialová

9th–11th May 2023, Křtiny

MENDEL UNIVERSITY IN BRNO

Czech Society of Landscape Engineers – ČSSI, z. s.,



and

Department of Landscape Management Faculty of Forestry and Wood Technology Mendel University in Brno



Public recreation and landscape protection – with environment hand in hand?

Proceedings of the 14th Conference

Editor: associate Professor Ing. Jitka Fialová, MSc., Ph.D.

9–11 May 2023 Křtiny Under the auspices

of prof. Dr. Ing. Jan Mareš, the Rector of Mendel University in Brno,

of prof. Dr. Ing. Libor Jankovský, the Dean of the Faculty of Forestry and Wood Technology, Mendel University in Brno,

of doc. Ing. Tomáš Vrška, Dr., the Director of Training Forest Enterprise Masaryk Forest Křtiny, Mendel University in Brno,

of Ing. Dalibor Šafařík, Ph.D., the Chief Executive Office, Forests of the Czech Republic,



of JUDr. Markéta Vaňková, the Mayor of the City of Brno,

BRNO

and of Mgr. Jan Grolich, the Governor of South Moravia,

south moravian region

in cooperation with Czech Bioclimatological Society, Nature Conservation Agency of the Czech Republic) and Partnerství, o.p.s.,

with the financial support of FS Bohemia Ltd.



The authors are responsible for the content of the article, publication ethics and the citation form.

All the articles were peer-reviewed.

© Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czechia

ISBN 978-80-7509-905-1 (print) ISBN 978-80-7509-904-4 (online ; pdf) ISSN 2336-6311 (print) ISSN 2336-632X (online ; pdf) https://doi.org/10.11118/978-80-7509-904-4

Open Access. This book is licensed under the terms of the Creative Commons Attribution 4.0 International License, CC-BY 4.0 (https://creativecommons.org/licenses/by/4.0/)



Contents

ADOLESCENTS' ENVIRONMENT Stanislav Azor, Mic	SMARTPHONE hal Marko, Štefan .	USAGE Adamčák	IN	ACTIVE	RECREATION	AND	NATURAL 9
ASSESSMENT OF OF SERBIA Martina Zeleňákova	EXTREME, LONG á, Milan Gocić, Har	G-TERM M	IETEC	DROLOGIC	AL DROUGHT II	N WEST atiana Sc	ERN PART)ľáková
AWARENESS OF Ľubomír Štrba, Br Sidor	SLOVAK TOURIST anislav Kršák, Len	rs ON POs ka Varcho	SSIBIL <i>lová,</i>	ITIES TO ' Michaela F	VISIT GEOSITES Podoláková, Silvi	S a Palgut	ová, Csaba 20
BANK STABILIZAT Miloslav Slezingr, I	TION – NON-TRAD Dana Košťálová	ITIONAL V	VAYS	OF USING	VEGETATION		24
BIOCULTURAL DI Ivo Machar, Helena	VERSITY: SACRAI a Kiliánová, Vilém F	L MONUMI Pechanec	ENTS	AS HABIT	ATS FOR BIRDS		27
CARAVANNING A Antonín Tůma	ND TRAMPING VE	RSUS CA	MPIN	G AND NA	TURE CONSERV	ATION	
CAUSES OF OVEI Emil Drápela	RCOMING OVERT	OURISM F	AILU	RE IN CZE	CHIA		
ECOTOURISM IN Petr Jelínek, Micha	AMAZONIAN ECU Il Hegar, Martin Mrl	ADOR – B kvička	OSQL	IE MEDICI	NAL PROJECT		
ERGONOMIC APF Eva Abramuszkino	PROACH IN TOUR vá Pavlíková, Osm	ISM FOR V an Nuri Öz	/ISITC dogan	RS WITH , Cihan Yi	SPECIAL NEEDS	8	45
EVALUATION OF Daniela Smetanova	THE LANDSCAPE	S POTEN	FIAL F	OR RECR	EATION		
EXPLORING THE Tímea Žolobaničov	RECREATIONAL F rá, Miroslav Čibik, F	POTENTIA Roberta Ště	L OF I épánko	URBAN GA ová	APS		57
EXPLORING WHIT Miroslav Čibik, Tím	TE SPACES ON UF lea Žolobaničová, F	RBAN MEN Roberta Ště	ITAL I ěpánko	MAPS ová			63
FOREST EDUCAT SOCIETY IN THE (Kamil Żołądek, Róż	TION AS THE BAS CELESTYNÓW FC ża Brytan, Artur Da	IS FOR COREST DIS	ONSC TRIC	IOUS USE	OF ALL FORE	ST FUN	CTIONS BY
GREEN AREAS AI OF RESIDENTS Edyta Rosion-Szer	ND NATURAL POT yńska	ENTIAL O	F THE	E POLISH (CITY OF CIESZY	'n in th	E OPINION
HEALTH VALUES Emilia Janeczko, N	OF FORESTS IN 1 Iałgorzata Woźnick	THE OPINI	ON O	F POLISH	RESIDENTS		79
HISTORICAL EV TOURISM INDUST Norie H. Palma, Ja	DLUTION OF FE RY GROWTH smin T. Gadian, Jo	STIVALS	IN G ano, F	GUIMARAS Revenlie G.	ISLAND: ITS Galapin, Petr Ku	IMPLIC	ATION TO 83

HOW HIGH ARE THE ECONOMIC BENEFITS OF OUTDOOR RECREATIONAL USE FROM THE NEWLY DESIGNATED LANDSCAPE PROTECTED AREA? Jan Melichar, Petr Pavelčík, David Zahradník, Marek Banaš, Radim Misiaček, Jana Hamanová, Martin Slaba, Viktorie Kováčová
HOW TO MEASURE AND USE NATURE-BASED RECREATION EFFECTS: EXAMPLE OF RESULTS FOR THE VLTAVA RIVER CASCADE Kateřina Mácová, Jan Melichar, Vojtěch Havlíček, Martin Heřmanovský, Filip Strnad, Pavel Fošumpaur, Karel Březina, Martin Hanel, Martin Horský, Tomáš Kašpar, Vojtěch Sýs
HOW TO SUPPORT CARBON SEQUESTRATION AND RECREATIONAL POTENTIAL AT THE SAME TIME Jan Deutscher, Jana Smolíková
CHANGE OF THE WAY THAT LANDSCAPE IS USED AND IT IS EFFECT ON THE RECREATIONAL AND TOURIST POTENTIAL Jan Szturc, Jan Prachowski, Jana Podhrázská ,, Petr Karásek,, Josef Kučera,,
IMPLEMENTATION OF GREEN INFRASTRUCTURE ELEMENTS TO IMPROVE RECREATION IN THE VILLAGE OF DRIENOV, SLOVAKIA Martina Zeleňáková, Natália Junáková
IMPLICATIONS OF THE NATURE OF FORESTRY AND WOOD ENTERPRISES IN LATVIA Dastan Bamwesigye, Ingus Grinbergs, Amanda Puzule, Tīna Ķikule
INFLUENCE OF RECREATIONAL ACTIVITIES ON THE DISTRIBUTION OF FOREST WILD BOAR ROOTING Jakub Drimaj, Marie Balková, Jakub Špoula, Jiří Kamler, Ondřej Mikulka, Radim Plhal, Miloslav Homolka
INNOVATIVE TECHNOLOGY OF SAPLINGS PLANTING FOR INCREASE TOURISM POTENTIAL OF THE LANDSCAPE Luboš Staněk, Ladislav Zvěřina, Radomír Ulrich
INTEGRATION OF NICHE AGRICULTURAL CROPS IN THE DEVELOPMENT OF ROMANIAN RURAL TOURISM. CASE STUDY: WALNUT CULTURE IN ROMANIA AFTER 1990 Constantin-Răzvan Oprea, Roxana Cuculici, Iulian Săndulache
INTELLIGENT DESTINATION GUIDE David Zejda, Martina Pásková130
LANDSCAPE – ARCHITECTURAL PROPOSAL OF JANDURA PARK IN CANBERRA, AUSTRALIA: EXPERIENCE OF BILATERAL COOPERATION BETWEEN TWO UNIVERSITIES Mária Bihuňová, Miroslav Čibik, Roberta Štěpánková, Attila Tóth
LANDSCAPE CHARACTER AND INTEGRATION OF MINING LAKES INTO THE LANDSCAPE - OPPORTUNITIES AND RISKS Hedvika Psotová
LANDSCAPE-ARCHITECTURAL SOLUTION AROUND THE RIVER VÁH IN THE CADASTRAL TERRITORY OF SEREĎ WITH AN EMPHASIS ON RECREATION Denis Bechera, Gabriel Kuczman, Miroslav Rusko144
LANDSCAPE-FRIENDLY METHOD OF FOUNDING WOODEN BUILDINGS FOR RECREATIONAL USE
ravia nulaskuva, jilka rialova
SHARED-USED RECREATIONAL TRAILS IN THE CZECH REPUBLIC Hana Hermová, Tomáš Kvasnička154

METEOROLOGICAL ACTIVITIES OF J. G. MENDEL AS PART OF THE TOUR OF THE AUGUSTINIAN ABBEY Jaroslav Rožnovský
MID-FIELD WOODLOTS AS A SUBSTITUTE FOR FORESTS IN AGRICULTURAL AREAS - THE IMPACT ON ENVIRONMENT AND TOURISM Beata Fortuna-Antoszkiewicz, Jan Łukaszkiewicz, Piotr Wisniewski
MONITORING THE MOVEMENT OF VISITORS IN THE TATRA NATIONAL PARK USING BATTERY-POWERED ONLINE COUNTERS Ivos Gajdorus
MOUNTAIN RESCUE SERVICE - INEVITABLE HELP AT RECREATIONAL AND SPORT ACTIVITIES IN MOUNTAINOUS AREAS IN SLOVAKIA Matúš Jakubis, Mariana Jakubisová
NON-WOOD FOREST PRODUCTS: "CULTURE" + "TRADITION" = "EDUCATIONAL POSSIBILITIES". DOES IT MAKE SENSE? Szczepan Kopeć, Paweł Staniszewski
OLDER ADULTS AS A TARGET GROUP OF USERS OF GREEN AREAS IN PROJECTS OF THE WARSAW CIVIC BUDGET Kinga Kimic, Paulina Polko
PLANNING THROUGH A GIS THE RECOVERY OF RURAL BUILDINGS FOR THE DEVELOPMENT OF NEW FORMS OF TOURISM HOSPITALITY <i>Pietro Picuno, Salvatore Margiotta188</i>
POSSIBILITIES AND ADVANTAGES OF INDIVIDUAL RECREATION IN THE TOPOĽČANY DISTRICT Regina Mišovičová, Zuzana Pucherová, Henrich Grežo,
POSSIBILITIES OF RECREATION IN HNILEC RIVER BASIN FROM CLIMATOLOGICAL POINT OF VIEW Patrik Nagy, Katarzyna Kubiak-Wójcicka , Miroslav Garaj , Milan Gocic3
POSSIBILITIES OF USING NEW TECHNOLOGIES IN CULTURAL TOURISM IN THE POST COVID ERA Kristýna Tuzová, Milada Šťastná202
PROBLEMS OF RURAL LANDSCAPE'S PROTECTION VS ANTHROPOPRESSURE AND RECREATION MOVEMENT - THE EXAMPLE OF THE NATURE RESERVE "STAWY RASZYŃSKIE" NEAR WARSAW Jan Łukaszkiewicz, Beata Fortuna-Antoszkiewicz
PUBLIC RECREATION AND TOURISM ARE ASPECTS THAT AFFECT NOT ONLY THE ENVIRONMENT David Brandejs, Pavel Klika
QUALITATIVE ASSESSMENT OF THE PREPAREDNESS AND POTENTIAL OF NATURE PROTECTED AREAS TO SUPPORT SUSTAINABLE TOURISM Radek Timoftej and Hana Brůhová Foltýnová
RECREATION IN CZECH LARGE PROTECTED AREAS: COUNTED AND SORTED Tomáš Janík
RECREATION LAND USE IN TERMS OF WATER PROTECTION Maria Hlinkova, Rastislav Fijko228

RECREATIONAL POTENTIAL OF RADOŠINKA MICROREGION: LANDSCAPE – ARCHITECTURAL PROPOSAL OF THE CYCLO ROUTE Mária Bihuňová, Branislav Králik
RECREATIONAL USE OF FOREST ROADS IN THE TERRITORY OF NATIONAL PARKS AND PROTECTED LANDSCAPE AREAS Roman Bystrický
REFORM OF THE CONSTRUCTION ADMINISTRATION IN RELATION TO THE PERMITTING OF BUILDINGS FOR RECREATION Alena Kliková
REVITALISATION OF DRAINED FOREST AREA Jana Marková, Petr Pelikán
REVITALIZATION OF THE PARK IN THE CENTER OF IVANKA PRI DUNAJI Gabriel Kuczman, Denis Bechera
RISK ASSSESSMENT ON GEODIVERSITY SITES Lucie Kubalíková, Eva Nováková, František Kuda, Karel Kirchner, Aleš Bajer, Marie Balková
RIVERS AS BACKBONES FOR URBAN AND PERIURBAN RECREATION – CASE STUDIES FROM KOŠICE AND PREŠOV, SLOVAKIA Juraj Illes, Katarina Kristianova
SMALL-SCALE INVASIVE INTERVENTIONS AS IMPULSES FOR THE REACTIVATION OF FORGOTTEN URBAN SPACES <i>Miroslav Čibik, Katarína Jankechová</i>
STUDY OF THE RELATIONSHIP OF MOISTURE AND COMPACTION ON THE MODULUS OF RESILIENCE OBTAINED BY CYCLIC CBR TESTING IN LOCAL SOILS FOR A QUALITY RURAL TOURISM Iñigo Garcia, Lenka Ševelová
THE "KAMIEŃ" EDUCATIONAL PAVILION IN WARSAW AS A PLACE OF PRO-ENVIRONMENTAL ACTIVATION OF THE URBAN COMMUNITY Kinga Kimic , Magdalena Wolska
THE ASSESSMENT OF ECOSYSTEM SERVICES IN TRNAVA (SLOVAKIA) AND SURROUNDING REGION Radovan Pondelík, Martin Zápotocký
THE CONCEPT OF SENSE OF PLACE IN ENVIRONMENTAL EDUCATION Dominik Rubáš, Tomáš Matějček, Tomáš Bendl
THE EFFECT OF GRASS STRIPS ON SOIL RETENTION AND EROSION REDUCTION IN AGRICULTURAL LANDSCAPE Petr Karásek, Josef Kučera, Michal Pochop
THE FIRST OFFICIAL FOREST MIND TRAIL IN THE CZECH REPUBLIC – KŘTINY ARBORETUM Jitka Fialová, Martina Holcová
THE HIPOROUTES IMPLEMENTATION OPTIONS FROM ALTERNATIVE MATERIALS Václav Mráz, Jiří Ježek , Karel Zlatuška , Vlastimil Nevrkla
THE IMPACT OF THE CREATION OF A RECREATIONAL AREA BY RECLAMATION OF A SURFACE MINE ON PROPERTY VOLUE Vítězslava Hlavinková, Martina Vařechová

THE IMPLEMENTATION OF GIS TOOLS FOR PLANNING THE DEVELOPMENT OF RURAL TOURISM ALONG THE NETWORK OF OLD SHEEP-TRACKS <i>Giuseppe Cillis, Dina Statuto, Pietro Picuno</i>
THE IMPORTANCE AND FUNCTIONS OF RIPARIAN STANDS OF THE RECREATIONAL WATER RESERVOIR POČÚVADLO IN ŠTIAVNICKÉ VRCHY Mariana Jakubisová , Matúš Jakubis
THE ISSUE OF GEO-EDUCATION ON NATURE TRAILS IN THE FIRST SLOVAK GEOPARK BANSKÁ ŠTIAVNICA Silvia Palgutová, Michaela Podoláková, Lenka Varcholová, Branislav Kršák, Ľubomír Štrba
THE ROAD FROM THE CITY TO THE FOREST. OR HOW FAR IS THE URBAN MAN FROM A FUNCTIONAL FOREST? Vilém Pechanec, Helena Kilianová, Ivo Machar
THE ROLE OF LAND CONSOLIDATION IN RURAL SPACE DEVELOPMENT Jana Konečná, Michal Pochop, Jana Podhrázská, Petr Karásek, Eva Nováková
THE ROLE OF WETLANDS IN FLOOD PROTECTION PROCESSES IN THE LANDSCAPE – CASE STUDY Marián Dobranský, Peter Bujanský, Gao Zhenjun
THE UNFINISHED HITLER'S MOTORWAY – A HERITAGE IN THE CONTEMPORARY LANDSCAPE Ivo Dostál, Marek Havlíček, Hana Skokanová
TRADITIONAL COPPICE MANAGEMENTS AT THE LANDSCAPE LEVEL TOGETHER WITH RECREATIONAL USE Barbora Uherková, Jan Kadavý , Zdeněk Adamec, Michal Friedl, Aleš Kučera, Robert Knott, Michal Kneifl, Jakub Drimaj
TRANSFORMATION OF GARDEN SETTLEMENTS INTO A RESIDENTIAL ZONE Sofie Pokorná, Vítězslava Hlavinková
UNDERGROUND SPACES IN BOSONOŽSKÝ HÁJEK NATURE RESERVE AND THEIR GEOEDUCATION IMPORTANCE Karel Kirchner, František Kuda, Vít Baldík, Lucie Kubalíková
URBAN AGRICULTURE – ECOSYSTEM AND CULTURAL FUNCTIONS OF ORCHARD VEGETATION Jan Winkler, Petra Martínez Barroso, Doubravka Kuříková, Helena Pluháčková, Aleksandra Nowysz
VALORIZATION OF AN OLD SHEEP TRAIL AS A NEW OPPORTUNITY FOR SUSTAINABLE PUBLIC RECREATION: A CASE STUDY IN SOUTHERN ITALY Dina Statuto, Giuseppe Cillis, Pietro Picuno
WHERE NATURE MEETS ADVENTURE: TOURIST ACTIVITIES AT DOBROGEI GORGE NATURE RESERVE, ROMANIA <i>Teodorescu Camelia, Szemkovics Laurentiu-Stefan, Dumitrascu Alina Viorica</i>
WHERE THE SQUARE MEETS THE STREAM: RE-DESIGNING THE RURAL SQUARE IN VEĽKÝ KÝR, SLOVAKIA Attila Tóth
WHICH INFLUENCE HAS DEFORESTATION ON TOURISTIC RECREATIONAL AREAS IN SUCEAVA COUNTY? Ana-Maria Ciobotaru

TRADITIONAL COPPICE MANAGEMENTS AT THE LANDSCAPE LEVEL ALONG WITH RECREATIONAL USE

Barbora Uherková¹, Jan Kadavý¹, Zdeněk Adamec¹, Michal Friedl², Aleš Kučera³, Robert Knott⁴, Michal Kneifl¹, Jakub Drimaj⁵

¹ Department of Forest Management and Applied Geoinformatics, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemědělská 3, 613 00 Brno, Czechia

² Department of Forest Botany, Dendrology and Geobiocenology, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemědělská 3, 613 00 Brno, Czechia

³ Department of Geology and Pedology, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemědělská 3, 613 00 Brno, Czechia

⁴ Department of Silviculture, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemědělská 3, 613 00 Brno, Czechia

⁵ Department of Forest Protection and Wildlife Management, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemědělská 3, 613 00 Brno, Czechia

https://doi.org/10.11118/978-80-7509-904-4-0346

Abstract

Our current goals in forestry involve safeguarding the land and environment, optimizing the forest's ability to absorb CO₂, maintaining the health and integrity of forest ecosystems, and preserving biodiversity and landscape variety. Ensuring the long-term productivity, socio-economic viability, and environmental functions of European forests and woodlands, including coppice woodlands, is crucial. Addressing these challenges can be effectively achieved by utilizing a combination of various management methods. The main focus of this study is on traditional forms of management such as coppicing, grazing, and litter raking. To facilitate this research, designated plots were established within the Training Forest Enterprise Masaryk Forest Křtiny. These plots, where sheep grazing and litter raking have been implemented since 2018, are exceptional and provide exemplary illustrations of diverse coppice management practices in Central European coppices.

Key words: standards, grazing, litter raking, biodiversity, recreational value

Introduction

Numerous cultural landscapes throughout Europe have been molded by centuries of coppice forest management. As the oldest form of forest management, coppicing has a long and rich history. Today, it is estimated that over 20 million hectares of forest in Europe are currently managed using coppice techniques. In the Czech Republic, coppicing was once a prevalent form of forest management, along with grazing and litter raking. However, this practice was eventually abandoned in the 19th century due, in part, to the shift towards using fossil fuels. Following World War II, coppicing was completely discontinued. During the early Communist period in the 1950s, many forestry researchers viewed coppicing as a "capitalist" approach that focused solely on maximizing wood production, even if it meant depleting soil nutrients and ultimately leading to soil degradation and decreased wood production capacity. Consequently, this led to the demise of the three traditional non-timber forest uses: coppicing, grazing, and litter raking, which were once widely practiced.

Today, with new information and perspectives, many European countries view coppicing differently. Coppice management practices are now recognized for their numerous benefits, including minimal soil damage during harvest (in the case of coppice with no standards), reduced need for weed management, physical protection of the site on sloping terrains, reduced risk of windthrow, and long-term soil stability. Additionally, where markets for coppice products exist, this management method can provide a financial return for landowners, particularly for those who own small forest properties.

Coppicing also provides crucial environmental benefits such as carbon sequestration, as well as conservation of water, soil, and biodiversity. This management practice supports a diverse range of forest species, and it also holds significant cultural and spiritual value. Furthermore, we should not overlook the recreational value of coppicing and grazing, which have become increasingly important in recent years. It is worth noting that tourist and recreational use of forests is one of the newest forest functions. Tourism and recreational activities in forests are highly popular in the Czech Republic. However, forest visitors are often conservative in their views and lack an understanding of new forest management practices. To address this, efforts are being made to educate both the public and professional communities about coppicing and to shift their perspectives towards this management

method. According to Stanturf (2015), reintroducing sheep grazing in forests can be viewed as a means of restoring a mosaic landscape, and it may enhance the appeal of specific forested areas. There is now a growing effort throughout Europe to maximize the benefits of this currently underutilized management system.

Coppicing is a valuable form of forest management that should be recognized as such. By promoting a diversity of forest management practices over space and time at the landscape level, biodiversity can be significantly enhanced, and heterogeneity can be restored to production forest landscapes. In fact, increasing the diversity of forest management practices is an effective approach to adapting to global change, according to Duflot et al. (2022).

The primary objective of this research is to assess the effects of traditional forest management practices on forest ecosystems, with a particular focus on biometric, pedological, and geobiocenological perspectives.

Materials and methods

The study conducted at the Training Forest Enterprise Masaryk Forest Křtiny involved the establishment of 15 plots, each measuring 40 × 30 m. These plots were established to evaluate the effects of traditional forest management practices and differ from each other based on the type of management treatment applied. The prevailing slope aspect and inclination at the study site is W-N and 5-10°, respectively, at an altitude of 275-325 m above sea level. Due to differing loess admixture (even more than 0.5 m of continuous layer on steeper western slope) in granodiorite bedrock, soil protection includes minimization of mineral soil exposure and continual plant cover. The set of research plots was established in 2017. Twelve of the plots were subjected to strong thinning, resulting in only 80 trees per hectare being retained as standards. Three plots were left without any further intervention as control plots. Dendrometrical measurements of the circumference of oak standards at breast height were carried out at the beginning and end of each growing season. The relative growth ratios (RGR) were computed using the formula provided by Cotillas et al. (2009), as shown below:

$$RGR = \frac{x_i - x_{i-1}}{x_{i-1}} * 100 \tag{1}$$

where x_i is the circumference at breast height in time *i* (end of the growing season) and x_{i-1} is the circumference at breast height in time *i*-1 (beginning of the growing season).

The study compared the RGRs of oak standards among five different treatments (control plots, plots with only coppice, and three treatments with coppice combined with litter raking, grazing, or both) over six years (with 2017 used as a control year before any treatments were applied). The ANOVA and post hoc multiple comparison Tukey tests were used to analyze and compare the RGRs.

Parametric ANOVA and post hoc multiple comparison Tuckey tests were used to analyze the concentrations of soil mineral nitrogen forms (ammonia NH_{4^+} and nitrates NO_{3^-}) in fresh samples collected during 2017-2021, as assessed by Kučera et al. (2013). The statistical analysis was conducted at a significance level of 0.05.

Geobiocoenological sampling, as described by Randuška et al. (1986), was conducted on all the plots. The sampling was carried out on permanent geobiocoenological plots measuring 15 × 15 m and mainly included the composition of plant species and their estimated cover.

Results and Discussion

The dendrometrical measurements of oak standards in the coppice plots revealed that there was an increase in the circumference growth, as expected. The maximum increment in circumference was observed four years after the start of the experiment, following which there was a decline in the increment (Figure 1). This could be attributed to the release of canopy and reduced competition resulting from the initial harvesting of trees. The results were consistent with the hypothesis proposed by Jones and Thomas (2004), which suggests that the diameter increment of trees released from competition is initially high but later becomes similar to the values before the release.



Fig. 1: Mean values (significance level α=0.05) of circumference relative growth ratio of oak standards in years 2017-2022 compared between different treatments. RGR (%) – relative growth ratio of oak standards, C – control plots, LR – litter raking plots, G – grazing plots (Adamec et al., 2022).

Pedologically, the results indicate an increase in ammonia nitrogen concentration during the first year after disturbance, as shown in Figure 2. The most intensive treatments, involving coppicing, litter raking, and grazing, resulted in concentrations more than five times higher compared to the control plot. From 2019 to 2021, the concentrations of ammonia nitrogen, as well as the differences among treatments, systematically decreased to the same level as in 2017. In contrast, nitrate nitrogen concentrations were consistently higher in 2018, with the highest concentration observed in the treatment combining all three management practices. Over time, nitrate concentrations continued to increase systematically. The concentrations of nitrate form result from the mineralization process following disturbance and inputs through faeces from grazing. Both forms can be utilized by nitrophilous plants, but nitrates are susceptible to leaching.



control plots; LR – litter raking; G – grazing (Adamec et al., 2022).

From a geobiocoenological perspective, our findings indicate that the reintroduction of traditional management practices resulted in a significant increase in the number of species in the herbaceous layer. Prior to the treatments, the number of species identified ranged from 7 to 26, with an average of 17 species per plot. Following the treatments, this number increased to between 61 and 117 species (excluding control plots) with an average of 86 species per plot in 2018, and to between 75 and 103 species with an average of 88 species per plot in 2019.

Figure 3 displays the average number of species per plot over time for each of the different treatments. The figure clearly shows that the combination of all three treatments (coppicing, grazing, and litter raking) resulted in the greatest increase in the number of species in the first year, followed by a decline in subsequent years. In contrast, the other treatments (coppicing alone, coppicing and raking, or coppicing and grazing) resulted in a lower increase in the number of species in the first year, but this increase continued in the following year. Two years after the treatment, the number of species, regardless of the treatment, started to converge. However, the most intensive treatment (i.e. the combination of coppicing, grazing, and raking) still maintained a higher species richness compared to the other treatments (see Fig. 3 and 4).



Fig 3: Growth of the mean number of species according to treatments between years 2017–2019. C = coppicing, R = litter raking, G = grazing.



Fig. 4: Change in the median number of species according to treatments between years 2017–2019. C = coppicing, R = litter raking, G = grazing.

Conclusion

This contribution presents the results of an experiment conducted at Training Forest Enterprise Masaryk Forest Kritiny in the southeastern part of the Czech Republic. The experiment involved applying coppicing, litter raking, and grazing to research plots. The results indicate that the released oak standards showed an increase in circumference increments during the first years of the experiment. The treatments also increased the concentrations of ammonia and nitrate forms of nitrogen, which can be beneficial to nitrophilous plants. Furthermore, the treatments increased the number of species in the herbaceous layer. However, further monitoring is needed to determine if the treatments differ from the control plots in the long term. Overall, these management practices can help increase the diversity of forest types at the micro-scale level and contribute to landscape diversity.

The ongoing project at TFE Masaryk Forest not only provides valuable insights into the effects of traditional forest management methods on forest ecosystems, but also offers a unique opportunity for visitors to witness practices that are no longer common in Czech forests due to legal restrictions. The project also aims to raise awareness about the importance of traditional forest management methods and their potential for enhancing forest biodiversity. With the research plots located near popular

tourist routes, it is expected that the project will attract a larger number of visitors to the area, thereby increasing public awareness about sustainable forest management practices.

References

Adamec, Z.; Friedl, M.; Kadavý, J.; Kneifl, M.; Knott, R.; Kománek, M.; Kostka, M.; Kučera, A.; Neubauer, Š.; Uherková, B.; Vichta, T. (2022). The influence of historical and modern forest managements on the condition of coppice – results in 2022. In SilvaNet – WoodNet 2022: Proceedings Abstracts of Student Scientific Conference. Brno: Mendelova univerzita v Brně, 2022, s. 12–13.

Cotillas, M.; Sabaté, S.; Gracia, C.; Espelta, J. M. (2009). Growth response of mixed mediterranean oak coppices to rainfall reduction: Could selective thinning have any influence on it? For. Eco. Manag. 2009, 258, 1677–1683.

Duflot, R.; Fahrig, L.; Mönkkönen, M. (2022). Management diversity begets biodiversity in production forest landscapes. Biol. Conserv. 2022, 268, 109514.

Jones, T.A.; Thomas, S.C. (2004). The time course of diameter increment responses to selection harvests in *Acer saccharum*. Can. J. For. Res. 2004, 34(7), 1525–1533.

Kučera, A.; Holík, L.; Marosz, K.; Martiník, A.; Vavříček, D. (2013). Changes in forms of available nitrogen and respiration in soil of beech forest as a reaction to a deforestation resulting from wind storm. Acta Univ. Agric. Silvic. Mendel. Brun. 2013, 61, 107–113.

Randuška, D.; Vorel, J.; Plíva K. (1986). Fytocenológia a lesnícka typológia. Bratislava: Príroda, 1986. Stanturf, J.A. (2015). Future landscapes: Opportunities and challenges. New For. 2015, 46, 615–644.

Acknowledgement

The research was supported by the Specific University Research Fund MENDELU, project No. IGA-LDF-23-TP-002.

Souhrn

Daný příspěvek vznikl v rámci probíhajícího projektu na ŠLP ML Křtiny, kde se zkoumá vliv řízené pastvy a hrabání opadu na stav pařezin. Dílčím cílem je separace těchto vlivů a kvantifikace jejich účinku na stav pařezin. Přičemž pařezina (les nízký) může být vnímána jako jeden z tvarů lesa se specifickým managementem, který vede k diverzifikaci způsobů hospodaření v čase a prostoru, a to na úrovni celé krajiny. Projektem sledované faktory (pastva a hrabání opadu a jejich vzájemné ovlivnění) v minulosti přispěly k negativnímu povědomí o pařezinách u odborné, ale i laické veřejnosti. V současné době se snažíme o nový náhled na tyto tradiční způsoby hospodaření.

Contact: Ing. Barbora Uherková, Ph.D. E-mail: xfedorov@mendelu.cz

Open Access. This article is licensed under the terms of the Creative Commons Attribution 4.0 International License, CC-BY 4.0 (<u>https://creativecommons.org/licenses/by/4.0/</u>)



Title: **Public recreation and landscape protection – with environment hand in hand?** Proceedings of the 14th Conference Editor of the proceeding: associate Professor Ing. Jitka Fialová, MSc., Ph.D. Publisher: Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czechia Print: Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czechia Edition: 1st Edition, 2023 No. of pages: 392 No. of copies: 75 ISBN 978-80-7509-905-1 (print) ISBN 978-80-7509-904-4 (online ; pdf) ISSN 2336-6311 (print) ISSN 2336-632X (online ; pdf) https://doi.org/10.11118/978-80-7509-904-4